II. *A Catalogue of the* 50 *Plants, from* Chelſea *Garden, preſented to the* Royal Society, *for the Year* 1724. *by the Company of Apothecaries, purſuant to the Direction of Sir* Hans Sloane *Bart. Pr. Coll. Med. S. P. V. Pr. By Mr.* Iſaac Rand, *Apothecary, F. R. S.*

101. GEranium lucidum, ſaxat ile. C.B. 318.
102. Geranium folio Malvæ rotundo. C. B. Ib. Majus, foliis in breviores & latas lacinias diviſis.
103. — Columbinum, vulgare, petalis florum bifidis, purpureis.
— Secundum *Camer. Epit.* 600.
104. — — Petalis florum bifidis albis. *An* Ger. Malacoides, flore albo. *H. Reg. Blæſ.*
105. — — Humile, flore cæruleo, minimo. *Raii Syn. Ed.* 3. p. 359.
— —Tenuius laciniatum. C. B. 318. *Prod.* 138.
106. — — Majus, flore minore, cæruleo. *Raii H.* 1058.
107. — — Maximum, diſſectis foliis. *Plot. Hiſt. Ox.*
108. — Orientale, columbinum, flore maximo, Aſphodeli radice. *T. Cor.* 20.
109. — Sanguineum, maximo flore. C. B. 318.
110. — — Cauliculis erectis, flore minore. *H. L. Bat.* 286.
111. — Hæmatodes, foliis majoribus, pallidioribus & altiùs inciſis, D. *Dale. Raii Syn* 219.

112. Geranium hæmatodes Lancaſtrenſe, flore eleganter ſtriato. *Raii Hiſt.* 1061.
113. —— Batrachoides. Gratia Dei, Germanorum. C. B. 318.
114. —— —— Montanum noſtras. *Raii Hiſt.* 1062.
115. —— —— Odoratum. C. B. 318.
116. —— Annuum minus Batrachoides, Bohemicum, purpureo-violaceum. *H. Ox.* p. 2. 511.
117. —— Phæum, flore reflexo I. B T. 3. 477.
118. —— Magnum, folio trifido I. B. T. 3. 478.
119. —— Romanum verſicolor, ſive ſtriatum. *Park. Parad.* 229.
120. —— Robertianum primum. C B. 319.
121. —— —— Rubens, flore albo.
122. —— Lucidum, ſaxatile, foliis Geranii Robertiani. *D. Sherard. Raii Syn.* 218.
123. —— Folio Althææ. C. B. 318.
124. —— —— —— Floribus ſaturatiùs rubentibus. *H. L. Bat.* 290.
125. —— Chium Vernum, Caryophyllatæ folio. *T. Cor.* 20.
126. —— Folio Alceæ, tenuiter laciniato; flore pentapetalo; purpuraſcente, ſemine tenui. *Boerh. Ind. alt.* 266.
127. —— Cicutæ folio, moſchatum. C. B. 319.
128. —— —— ——, Minus & ſupinum C. B. 319.
129. —— Arvenſe, album. *Tabern. Ic.* 57.
130. —— Myrrhinum, tenuifolium, amplo flore purpureo. *Barrel.* ob. 55.
131. —— Latifolium, longiſſimâ acu. C. B. 319.
132. —— Cicutæ folio, acu longiſſimâ. C. B. ib.
133. —— Acu longiſſimâ, Cicutæ folio tenuiùs laciniato.
—— Cicutæ folio, viſcoſum, acu longiſſima. D. *Juſ. ſieu.*

134.

134. —— Triſte *Cornuti.* 109.

135. —— Æthiopicum, noctu olens, radice tuberoſa, foliis Myrrhidis, anguſtioribus. *Breyn. Cent.*

136. —— Africanum, Coriandri folio, floribus incarnatis, minus. *H. L. Bat.* 279.

137. —— Africanum fruteſcens, Chelidonii folio, petalis florum anguſtis, albidis; carnoſo caudice.

138. —— —— Noctu olens, tuberoſum & nodoſum, Aquilegii foliis. *H. L. Bat.* 284.

139. —— —— Betonicæ folio, procumbens, floribus parvis, eleganter variegatis. *Pluk. Alm.* 169.

140. —— —— Foliis inferioribus Aſari, ſuperioribus Staphydis agriæ, maculatis, ſplendentibus, & Acetoſæ ſapore. *Com. Præl.* 52.

141. —— Africanum fruteſcens, folio craſſo & glauco, Acetoſæ ſapore. *Com. Præl.* 54.

142. —— —— Alchimillæ hirſuto folio, floribus albidis. *H. L. Bat.* 282.

143. —— —— Folio Malvæ craſſo, molli, odoratiſſimo, floſculo pentapetalo, albo. *Boerh. Ind. alt.* 263.

144. —— —— Fruteſcens, Malvæ folio laciniato, odorato. *H. L. Bat.* 277.

145. —— —— —— —— —— —— Inſtar Meliſſæ, flore purpuraſcente. *Boerh. Ind.* 110.

146. —— —— Arboreſcens, Malvæ folio mucronato, petalis florum inferioribus vix conſpicuis.

147. —— —— —— Alchimillæ hirſuto folio, floribus rubicundis. Com. præl. 51.

148. —— —— —— Folio Malvæ plano, lucido, glabro; flore Kermeſino, Di van Leur. *Boerh. Ind.* 109.

149. —— —— —— Ibiſci folio rotundo; Carlinæ odore. *H. L. Bat.* 274.

150. —— —— ——, —— —— Anguloſo; floribus amplis purpureis.

—— —— Fruteſcens, Alchimillæ folio, hirſuto, cum fimbriis purpureis. *Boerh. Ind.* 110.

III. *Obſer-*